



After surgery instructions to parents



Give Clear liquids for the first 3-4 hours after surgery (Ex: water, broth, flavored water, clear juice, etc.)



For the remaining of the day, soft diet is recommended (soup, jello, apple sauce, etc.)



Your child may be drowsy for some time after the procedure so limited activity is recommended for the remainder of the day.



If your child wants to sleep. Position him/her on his/her side with head supported and chin up.



Tylenol or Motrin as needed for pain



Low grade fever (temperature to 100.5°) is common.

- Treat with Tylenol or Motrin as needed (follow the instructions on the bottle for doing)
- Dehydration may cause a slight increase in temperature- encourage clear fluids.
- If a higher fever develops or the fever persist, call University Surgery Center (209) 580-3400, _____, or _____



Nausea and vomiting are occasional side effects. If vomiting occurs, clear the material from the mouth. Be sure that breathing is normal. If vomiting persists for 20-40 minutes, contact our office at (209) 383-7804.



Use moistened gauze to clean the teeth and mouth for 1-2 days after surgery.



Regular brushing and flossing by ____ days after surgery.



Follow up dental appointment at Merced Pediatric Dentistry, if necessary.



If you have any questions regarding dental treatment feel free to contact our office at (209) 383-7804 or Dr. _____

Anesthesiologist:

Dr. _____

Merced Pediatric Dentistry
Dr. Moran, and Dr. Lee
113 W. El Portal Dr. Suite B, Merced CA 95348
www.mercedpediatricdentistry.com

Office use only